

Strangeness production in d+Au collisions at $\sqrt{s_{NN}} = 200$ GeV in STAR

Xianglei Zhu^a for the STAR collaboration

^aDepartment of Engineering Physics, Tsinghua University, Beijing 100084, China

Abstract

We report on the measurements of strange hadron (K_S^0 , Λ , Ξ) production in the most central (0-20%) d+Au collisions at $\sqrt{s_{NN}} = 200$ GeV in STAR. Significant strangeness enhancement is observed in the most central d+Au collisions, especially for multi-strange hyperons (Ξ). The nuclear modification factor (R_{dAu}) of identified particles shows possible particle type – baryon (protons, Λ), meson (π, K_S^0, ϕ) – dependence at intermediate p_T (2 - 3.5 GeV/c), indicating that the final state may have an impact on the Cronin effect.

1. Introduction

Enhancement of strange hadrons' yield in high energy nuclear collisions relative to that in $p + p$ collisions at the same energy has long been regarded as one of the signatures of Quark-Gluon Plasma (QGP) production in these collisions [1]. Supposedly, it will be more efficient to create $s\bar{s}$ pair in the QGP than in hadronic matter. In experiments, substantial strangeness enhancement has been observed in Au+Au and Cu+Cu collisions in STAR, especially for multi-strange hyperons [2, 3, 4]. According to canonical statistical model, strangeness enhancement in nuclear collisions can also be due to canonical suppression of strangeness production in $p + p$ collisions [5]. Recently, STAR's ϕ meson data [6, 7] shows an above unity enhancement in heavy ion collisions at RHIC, which seems favor the approach of a fully equilibrated core plus hadronic corona [8]. The study of strangeness production in d+Au collisions will connect A+A and $p + p$ data, thus leading to better understanding of strangeness enhancement in nuclear collisions.

The Cronin effect, which is the enhancement of hadron spectra at intermediate p_T in p+A collisions with increasing nuclear size, was first seen in the nuclear modification factor in low energy p+A collisions [9]. At RHIC energy, the nuclear modification factor in d+Au collisions for pions and protons [10] show that the Cronin effect exists for both particle species. The conventional models of the Cronin effect attribute the change of p_T spectra in p+A to multiple parton scattering in the initial stage before hard parton scattering, which will broaden the outgoing parton/hadron p_T distribution [11]. Recently, the coalescence/recombination model, which describes the hadronization of partons in deconfined matter and reproduces well many features of hadron spectra, v_2 and R_{CP} in Au+Au collisions at RHIC, has been applied to d+Au collisions to describe the hadron spectra and the Cronin effect [12]. Moreover, the recombination model predicts particle type (baryon/meson) dependence of the Cronin effect, as the particle type dependence of R_{CP} or v_2 at intermediate p_T in Au+Au collisions at RHIC. Therefore, the measurements of the Cronin effect for more particle species are helpful to understand the mechanism of the Cronin effect.

In STAR, strange hadrons are reconstructed with the topology of their weak decay channels, $K_S^0 \rightarrow \pi^+\pi^-$ (69.2% branching ratio), $\Lambda \rightarrow p\pi$ (63.9% branching ratio) and $\Xi \rightarrow \Lambda\pi$ (99.9% branching ratio)[13, 14]. This analysis is based on about 32 million minimum bias d+Au collision events from STAR's 2008 run. For the time being, only the most central events (0-20%) are well defined ($N_{part} = 15.2$ and $N_{bin} = 14.6$ according to Glauber model). In this paper, we present the results for the most central 20% events only.

2. Results




Figure 1: Transverse momentum spectra of K_S^0 , $\Lambda(\bar{\Lambda})$, $\Xi^-(\bar{\Xi}^+)$ at $|\eta| < 1.0$ from the most central (0-20%) d+Au collisions at $\sqrt{s_{NN}} = 200$ GeV. The $\Lambda(\bar{\Lambda})$ spectra were corrected by the weak decay from Ξ and Ξ^0 . The curves are Levy function fits to the spectra (solid, dot-dashed and dashed are for K_S^0 , Λ and Ξ^- respectively). Errors are statistical only.

After correcting the raw spectra by the acceptance, reconstruction and vertex finding efficiencies, which is estimated with an embedding method, we get the p_T spectra of K_S^0 , $\Lambda(\bar{\Lambda})$, $\Xi^-(\bar{\Xi}^+)$ shown in Fig. 1 at mid-rapidity ($|\eta| < 1.0$) for the most central d+Au collisions (0-20%). The $\Lambda(\bar{\Lambda})$ spectra have been corrected by the feed-down contributions from multi-strange baryon (Ξ and Ξ^0) weak decays. We assume Ξ^0 make the same contribution to Λ as the Ξ 's. The contribution from Ω is neglected in this analysis. The total feed-down contribution from Ξ 's is on the order of 23% to the inclusive Λ yield. The spectra can be well fitted with a Levy function [15], as shown in Fig. 1,

$$\frac{1}{2\pi p_T} \frac{dN^2}{dp_T dy} = dN/dy \frac{(n-1)(n-2)}{2\pi n T (nT + m(n-2))} \left(1 + \frac{\sqrt{p_T^2 + m^2} - m}{nT}\right)^{-n}. \quad (1)$$

In the Levy function, the soft component at lower p_T in the spectra is controlled by the parameter T , and the power law component at higher p_T is controlled by the index n . The integrated yield (dN/dy), dominated by the low p_T production, can be extracted from the fit. The percentage of dN/dy from extrapolation at low p_T is 10% for K_S^0 , 18% for Λ and 29% for Ξ .

Fig. 2 shows the enhancement factor ($dN/dy/N_{part}$ relative to that in $p + p$ collisions) as a function of N_{part} for K_S^0 (left), Λ , Ξ^- (middle) and $\bar{\Lambda}$, Ξ^+ (right) from d+Au (open symbols), Cu+Cu (grey) and Au+Au (black) collisions at $\sqrt{s_{NN}} = 200$ GeV. The details and interpretation of the Cu+Cu and Au+Au data can be found in [4]. Fig. 2 shows that there is also significant strangeness enhancement in the most central d+Au collisions, especially for multi-strange hyperon Ξ . The enhancement factor is almost the same for K_S^0 and Λ , and much larger for Ξ , which shows that the enhancement factor is proportional to the strangeness content. This strangeness content dependence of the enhancement factor is consistent with the predictions from the canonical statistical model [5].




Figure 2: Strangeness enhancement factor ($dN/dy/N_{part}$ relative to that in $p + p$ collisions) for K_S^0 (left), Λ , Ξ^- (middle) and $\bar{\Lambda}$, Ξ^+ (right) from d+Au (open symbols), Cu+Cu (grey) and Au+Au (black) collisions at $\sqrt{s_{NN}} = 200$ GeV. For d+Au data, errors are statistical only. The details of the Cu+Cu and Au+Au data can be found in [4]

Fig. 3 shows the nuclear modification factor (R_{dAu}) of strange hadrons (K_S^0 , Λ , $\bar{\Lambda}$, ϕ), as well as those of protons and pions which are shown as cyan and meshed brown bands respectively. STAR ϕ data is from [17]. The R_{dAu} for each identified particle is similar and lower than unity in the low p_T region. When $p_T > 1$ GeV/c, the R_{dAu} values of all particles are above unity, while baryons (Λ and proton) R_{dAu} rises faster than mesons. At intermediate p_T (2 – 3.5 GeV/c), R_{dAu} is grouped into two bands: baryons at about 1.81 ± 0.05 and mesons around 1.28 ± 0.05 . It should be noted here that ϕ mass is close to Λ mass, but its R_{dAu} is closer to K_S^0 's. This indicates the possible particle type dependence of R_{dAu} instead of particle mass dependence. This possible particle type (baryon/meson) dependence of R_{dAu} at intermediate p_T is similar to R_{CP} measurements in Au+Au collisions at the same energy [16]. When p_T increases beyond 3.5 GeV/c (~ 3 GeV/c for ϕ meson), the errors in strange hadrons' R_{dAu} become too large to draw any conclusion.

3. Summary

In summary, we presented the transverse momentum spectra, strangeness enhancement factor and nuclear modification factor R_{dAu} for K_S^0 , Λ and Ξ from the most central $\sqrt{s_{NN}} = 200$ GeV d+Au collisions at RHIC. Data shows significant strangeness enhancement in the most central d+Au collisions, especially for multi-strange hyperon Ξ . The enhancement factor is almost the same for K_S^0 and Λ and much higher for Ξ , which seems consistent with the predictions from




Figure 3: R_{dAu} of strange hadrons (K_S^0 , $\Lambda + \bar{\Lambda}$, ϕ) at mid-rapidity from $\sqrt{s_{NN}} = 200$ GeV d+Au collisions. R_{dAu} of protons and pions are shown as cyan and meshed brown bands respectively [10]. STAR ϕ data is from [17]. Errors are statistical only for pions, protons, K_S^0 and Λ . The left grey band is the 18% normalization error.

the canonical statistical model. The R_{dAu} data shows Cronin enhancement at intermediate p_T and has possible particle type (baryon/meson) dependence. This particular behavior can be naturally explained in the coalescence/recombination model, which shows that the hadronization process in the final state may have an impact on the Cronin effect.

Acknowledgments

This work was in part supported by China Postdoctoral Science Foundation.

References

- [1] J. Rafelski and B. Müller, Phys. Rev. Lett. **48**, 1066 (1982)
- [2] B. I. Abelev *et al.*, Phys. Rev. C **77**, 044908 (2008)
- [3] J. Adams *et al.*, Phys. Rev. Lett. **98**, 062301 (2007)
- [4] A. Timmins, *these proceedings*; X. Wang (STAR Collaboration), J. Phys. G **35**, 104074 (2008)
- [5] K. Redlich and A. Tounsi, Eur. Phys. J. C **24**, 589 (2002)
- [6] N. Xu (STAR Collaboration), J. Phys. G **35**, 044031 (2008)
- [7] B. I. Abelev *et al.*, Phys. Lett. B **673**, 183 (2009)
- [8] F. Becattini and J. Manninen, Phys. Lett. B **673**, 19 (2009)
- [9] J.W. Cronin *et al.*, Phys. Rev. D **11**, 3105 (1975)
- [10] J. Adams *et al.*, Phys. Lett. B **616**, 8 (2005); *ibid*, Phys. Lett. B **637**, 161 (2006)
- [11] A. Accardi, arXiv:hep-ph/0212148
- [12] R. C. Hwa and C. B. Yang, Phys. Rev. Lett. **93**, 082302 (2004); R. C. Hwa and C. B. Yang, Phys. Rev. C **70**, 037901 (2004)
- [13] C. Adler *et al.*, Phys. Rev. Lett. **89**, 092301 (2002)
- [14] J. Adams *et al.*, Phys. Rev. Lett. **92**, 182301 (2004)
- [15] G. Wilk and Z. Włodarczyk, Phys. Rev. Lett. **84**, 2770 (2000)
- [16] J. Adams *et al.*, Nucl. Phys. A **757**, 102 (2005)
- [17] B. I. Abelev *et al.*, Phys. Rev. C **79**, 64903 (2009)